

The massive stellar content in NGC 604 and its evolutionary state.

Rosa M. González Delgado & Enrique Pérez

Instituto de Astrofísica de Andalucía (CSIC). Apdo. 3004, 18080 Granada, Spain

Accepted 1999. Received 1999; in original form 1999

ABSTRACT

This paper analyzes the integrated ultraviolet spectra taken with the *International Ultraviolet Explorer (IUE)* and optical ground-based spectra taken with the William Herschel Telescope (WHT) of the giant H II region NGC 604. These data are complemented with ultraviolet (WFPC2 through F170W) and H α (WFPC2 through F656N) images retrieved from the *Hubble Space Telescope (HST)* archive.

The ultraviolet resonance wind stellar lines, the nebular optical emission lines and the higher order terms of the Balmer series and HeI absorption lines detected in the spectra of NGC 604 are interpreted using evolutionary models optimized for young star forming regions. The evolutionary state and the massive stellar content of the region is derived in a self-consistent way.

The three techniques applied suggest that the central ionizing cluster in NGC 604 is very young, $\simeq 3$ Myr old, and that the stars in the cluster were formed in an instantaneous burst following a Salpeter ($\alpha=2.35$) or flatter ($\alpha=1.5$) initial mass function (IMF), having stars more massive than $80 M_{\odot}$. The stellar cluster is able to provide most of the ionizing photons needed to photoionize the whole nebula, and the wind power to form the central shell structure where the cluster core is located. The stellar cluster is affected by an extinction similar to the average extinction that affects the ionized gas. The estimated number of massive stars in the cluster is also in agreement with that derived from previous studies based on the detection of individual stars. The results that we present here support the use of these techniques for the interpretation of the integrated light of more distant star forming regions.

Key words: techniques: spectroscopic - galaxies: stellar clusters - ultraviolet: stars - ISM: H II regions - ISM: individual: NGC 604.

1 INTRODUCTION

Giant extragalactic H II regions (RH IIs) are amongst the brightest objects in galaxies. RH IIs have been studied extensively because they are the best indicators of the conditions that lead to massive star formation, and they show the cloud properties immediately after the moment when stars form. Equally important, they are favourable places where to measure the chemical abundances and star formation rates in galaxies, and hence can be used to trace the chemical and star formation history of the universe, when observed in galaxies at different redshift (Madau et al. 1996; Kobulnicky & Zaritsky 1999).

RH IIs are characterized to have a size larger than 100 pc and H α luminosity brighter than 10^{39} erg s $^{-1}$ (Kennicutt 1984). Therefore, the nebula requires an ionizing photon luminosity larger than 10^{51} s $^{-1}$; this is provided by a stellar cluster that contains more than 100 young massive

stars. These characteristics are very similar to those of starburst galaxies. However, they are less luminous than prototypical starbursts, and thus they are referred to as mini-starbursts (Walborn 1991). Like starbursts, RH IIs show a nebular emission-line spectrum at optical wavelengths, and an absorption-line spectrum at wavelengths shorter than the Balmer jump (Leitherer 1997; Rosa, Joubert & Benvenuti 1984). This spectral morphology reflects the fact that RH IIs are powered by massive stars. These stars emit photons with energies of tens of eV which are absorbed and re-emitted in their stellar winds, producing ultraviolet resonance transitions. However, the stellar wind is optically thin to most of the ultraviolet photons, that can travel tens of parsec from the star before they are absorbed and photoionize the surrounding interstellar medium. Subsequently, this ionized gas cools down via an emission spectrum. This spectral dichotomy picture allows to derive the stellar content and the evolutionary state of the cluster, through the anal-

ysis of the ultraviolet (e.g. Vacca et al. 1995; Leitherer et al. 1996), or the optical light (e.g. Cid Fernandes et al. 1992; García-Vargas & Díaz 1994; Stasińska & Leitherer 1996) using evolutionary synthesis and photoionization models. The two techniques have been applied to the prototypical starburst nucleus NGC 7714, giving similar results (González Delgado et al. 1999a).

However, around the Balmer jump the spectra of starbursts (González Delgado et al. 1998) and some H II regions (Terlevich et al. 1996) show the higher order terms of the Balmer series and He I lines in absorption, formed in the photospheres of massive stars. These stellar lines can be detected in absorption because the strength of the gaseous Balmer lines in emission decreases rapidly with decreasing wavelength, whereas the equivalent width of the stellar absorption lines is almost constant with wavelength (González Delgado, Leitherer & Heckman 1999b). Evolutionary synthesis models that predict the profiles of the higher order terms of the Balmer series and He I lines in absorption can also be used to estimate the evolutionary state of the stellar cluster in H II regions.

The second more luminous and brightest H II region in the Local Group of galaxies, after 30 Dor in the LMC, is NGC 604 in M 33. Its distance, 840 kpc (Freedman et al. 1991), allows detailed studies of the individual stars, and also to obtain integrated properties. Thus, NGC 604 is an excellent laboratory in which to explore questions about the effect of star formation, the slope and upper mass limit of the IMF, and to test the consistency between the three techniques described above, that allow to determine the stellar content and the evolutionary state of the ionizing stellar cluster from the spatially integrated ultraviolet and optical spectra of the H II region.

NGC 604 has been intensively studied in the past (e.g. Peimbert 1970; Israel & van der Kruit 1974; Smith 1975; Hawley & Grandi 1977; Kwitter & Aller 1981; Conti & Massey 1981; Rosa & D’Odorico 1982; Viallefond & Goss 1986). A precise determination of the chemical composition of the gas was made by Díaz et al. (1987); they give an oxygen abundance $12+\log(\text{O}/\text{H})=8.5$. VLA observations indicate that the gas is very tenuous (average electron density, $rms\ N_e$, between a few and $\sim 100\text{ cm}^{-3}$), and not very dusty, with a mean visual extinction $A_V \sim 0.5$ mag (Churchwell & Goss 1999). The morphology of the ionized gas is very complex, showing many filaments and shell structures that are expanding (e.g. Hippelein & Fried 1984; Rosa & Solf 1984; Clayton 1988; Sabalish et al. 1995; Muñoz-Tuñón et al. 1995; Yang et al. 1995; Medina Tanco et al. 1997). This morphology is a consequence of the violent star formation activity in NGC 604. *HST* imaging photometry has revealed that the stellar cluster is resolved into ~ 200 massive stars in an area of $\sim 10000\text{ pc}^2$ (Drissen, Moffat & Shara 1993; Hunter et al. 1996). Evidences that the stellar cluster is evolved come from the detection of Wolf-Rayet stars (Conti & Massey 1981; D’Odorico & Rosa 1981; Drissen, Moffat & Shara 1990, 1993), one supernova remnant (D’Odorico et al. 1980), and one candidate to red supergiant (Terlevich et al. 1996). The existence of these stars as members of the stellar cluster of NGC 604 suggests that the age of the region is in the range 3-5 Myr.

This paper presents spatially integrated ultraviolet and optical spectra of NGC 604. The goal is to derive the mas-

Figure 1. *HST*+WFPC2 (F170W) image of NGC 604 at ultraviolet wavelengths. The scale is $0.046\text{ arcsec pixel}^{-1}$, equivalent to $0.19\text{ pc pixel}^{-1}$. The rectangles mark the positions of the *IUE* apertures. North is up and East to the left.

Figure 2. *HST*+WFPC2 (F653N) image in the $\text{H}\alpha$ emission line (grayscale). The ultraviolet emission (F170W) is overlaid in white contours. The scale is $0.1\text{ arcsec pixel}^{-1}$. The lines indicate the slits of the ground-based optical spectroscopy. Note that the core of the cluster is within a nebular hole of 20 pc radius. North is up and East to the left.

sive stellar content and the evolutionary state of the ionizing cluster, by means of evolutionary synthesis and photoionization models applied to the ultraviolet resonance wind stellar lines, to the nebular emission lines and to the higher-order terms of the Balmer series and He I lines in absorption. The consistency between the results obtained with the three techniques, and the similarity with the results obtained from studies based on the detection of individual stars, strengthens the reliability and power of these techniques when they are applied to determine the stellar content of more distant star-forming regions. Section 2 presents the observations. Section 3 describes the photoionization models that fit the emission-line spectrum of the nebula. In section 4, the higher-order terms of the Balmer and He I lines in absorption are analyzed. Evolutionary synthesis models of the ultraviolet resonance wind stellar lines are in section 5. In section 6, the massive stellar content is derived. The summary and conclusions are in section 7.

2 OBSERVATIONS

2.1 *HST* images

NGC 604 has been intensively observed with the WFPC2 camera onboard the *HST*. For the purpose of this work, we have retrieved *HST* archive images at the ultraviolet (F170W filter) and optical $\text{H}\alpha$ (filter F656N) wavelengths. Two F170W exposures of 350 s each and two F656N exposures of 1000 s each were combined to produce one final ultraviolet (Fig. 1) and one $\text{H}\alpha$ image (Fig. 2). The nebula was centred in the PC camera, therefore the ionizing cluster is sampled at $0.046\text{ arcsec pixel}^{-1}$, corresponding to $0.19\text{ pc pixel}^{-1}$ (at a distance of 840 kpc, 1 arcsec corresponds to 4.1 pc). The ultraviolet emission is spread in the inner $20\times 20\text{ arcsec}$. The stellar density looks much lower than that observed in super star clusters. Surface brightness photometry indicates that 75 percent of the ultraviolet flux is produced by the core of the stellar cluster, that is contained within a central nebular hole of diameter $\sim 15\text{ arcsec}$; we shall refer to this as the centre of the nebula. The $\text{H}\alpha$ image shows many filaments and shell structures that extend out to $\sim 200\text{ pc}$ from the centre of the nebula. However, the brightest parts of the nebula trace almost a ring structure at a distance of 40 pc from the centre.

The total ultraviolet flux per unit wavelength in the F170W image is $1.9 \times 10^{-13}\text{ erg s}^{-1}\text{ cm}^{-2}\text{ \AA}^{-1}$. The $\text{H}\alpha$ flux is $4.0 \times 10^{-11}\text{ erg s}^{-1}\text{ cm}^{-2}$, in good agreement with the value measured by Churchwell & Goss (1999).

Table 2. Input parameters to CLOUDY.

12+log(O/H)	8.51
log (He/H)	-1.08
log (N/H)	-4.65
log (Ne/H)	-4.27
log (S/H)	-5.05
log Q	51.54 (s ⁻¹)
Inner Radius	20 pc
Ne	30, 100 cm ⁻³

of the gas to the ionizing source. For a spherical geometry, the average U is proportional to $(\phi^2 N_e Q)^{1/3}$.

The chemical abundances are fixed to those derived by Díaz et al. (1987). We scale the chemical composition of the gas to the value of the oxygen abundance, 12+log(O/H)=8.5, except the abundance of N, Ne, S and He, for which we take the values derived by Díaz et al. (1987; see Table 2). The electron density derived by Díaz et al. (1987) is of the order of 100 cm⁻³. Our measurements of the [S II]6717/6731 ratio indicate the low-density limit, with values below 100 cm⁻³ §. We have computed constant density models at two values of the electron density, 100 cm⁻³ and 30 cm⁻³. The latter value comes from the filling factor and an *rms* electron density of 10 cm⁻³, derived from the H α image. Also, an electron density of 30 cm⁻³ predicts a [S II]6717/6731 ratio which is in better agreement with the observed value.

3.2 The radiation field

The radiation field used as input to CLOUDY is the spectral energy distribution generated by the evolutionary synthesis code developed by Leitherer and collaborators (Leitherer et al. 1999). The code includes the new set of stellar evolutionary models of the Geneva group (Schaller et al. 1992; Schaerer et al. 1993a,b; Charbonnel et al. 1993; Meynet et al. 1994), and the stellar atmospheres grid compiled by Lejeune et al. (1997), supplemented by the expanding spherical non-LTE models of Schmutz, Leitherer & Gruenwald (1992). The latter stellar atmosphere models are applied to stars with very strong stellar winds. The spectral energy distribution was generated using the Z=0.008 (Z $\odot$ /2.5) metallicity tracks, assuming that stars and gas have the same metallicity. We assume two different star formation scenarios: instantaneous burst and continuous star formation at a constant rate. Different models are also computed with different assumptions about the IMF. The slope is Salpeter ($\alpha=2.35$), flatter ($\alpha=1.5$), or steeper ($\alpha=3.3$). The upper mass limit cut-off is set to 30, 60, 80 or 120 M $\odot$. The lower mass limit is fixed to 1 M $\odot$. However, photoionization models are not influenced by the lower limit mass cut-off if it is below 10 M $\odot$. The spectral energy distribution is normalized to an ionizing photon luminosity of log Q = 51.54 s⁻¹. This value is derived from the total H α flux measured in the image, after correcting by the mean extinction, $A_V = 0.5$ mag, derived by Churchwell & Goss (1999). We will check

below the compatibility of this normalization with the mass of the stellar cluster derived from the ultraviolet continuum luminosity.

3.3 The observational constraints

The emission lines were measured in our scanned spectrum of the nebula. The observed ratios have been corrected by the mean extinction, $c(H\beta)=0.22$, derived by Churchwell & Goss (1999). This value is in agreement with the reddening derived from the H γ /H β ratio measured in the scanned spectrum, when this ratio is corrected by the underlying stellar absorption. This extinction is lower than the values derived by Díaz et al. (1987), $c(H\beta)=0.3-0.4$; however, the extinction changes across NGC 604 (Maíz-Apellániz 1999). The de-reddened emission line ratios to be fitted by the photoionization models are included in Table 3. This table also includes the emission line ratios derived by Díaz et al. (1987) in several parts of the nebula, and those from an integrated spectrum in Maíz-Apellániz (1999). The range of values for each of these emission line ratios defines the tolerance that we accept for the difference between the observed and the predicted values from the models.

Following Stasińska & Schaerer (1999), we try to fit the strength of the emission lines, and also emission-line ratios indicative of the electron temperature ([O III]4363/5007), electron density ([S II]6717/6732) and ionization structure ([O III]5007/[O II]3727, [S III]9069/[S II]6716+6732). Another important observational constraint is the radius of the nebula. The H α image shows that the flux extends out to 200 pc of the center of the region; however, more than 90, 75 and 50 percent of the total flux is within the inner 140 pc, 100 pc and 60 pc radii, respectively (Fig. 3). Our determination of the core radius of the region is 80 pc.

3.4 Model results

We start by fixing the filling factor and the electron density by means of fitting the line ratios [S II]6717+6731/H β and [S II]6717/6731, and the radius of the nebula. [S II]6717+6731/H β is a good calibrator of the ionization parameter, because this ratio does not depend much on the IMF assumptions, on the star formation law or on the evolutionary state of the cluster (González Delgado et al. 1999a). Instead, this ratio depends on the filling factor, and for fixed geometry, ionizing photon luminosity and electron density, it depends on the ionization parameter ¶. The observed ratio, [S II]6717+6731/H $\beta \simeq 0.4$, indicates a filling factor $\phi \simeq 0.1$ (Fig. 6). Therefore, a large fraction of the volume of the region is filled with ionized gas; however, this gas is very tenuous because the *rms* electron density measured in the H α image is $\simeq 10$ cm⁻³, and the electron density in the S⁺ zone, derived from the definition of filling factor ($\phi = (N_e(rms)/N_e)^2$), is $N_e \simeq 30$ cm⁻³. The observed ratio [S II]6717/6731=1.4, indicates that the electron density is ≤ 100 cm⁻³. Models with a filling factor of 0.1 and $N_e = 30$

§ Notice, however, that the atomic parameters for the sulfur have changed in the intervening time

¶ Nonetheless, we should always be cautious when using sulfur dependent diagnostics, given the uncertainties about the atomic parameters and its not so well known ionization; also this calibration relies on the assumption of an ionization bounded nebula.




Figure 7. Predicted emission-line ratios, η parameter and the electron temperature as a function of the cluster age, for an instantaneous burst (filled symbols) and continuous star formation (open symbols). The models assume that the mass of the cluster is distributed following a Salpeter IMF between 1 and 120 $M_{\odot}$. The filling factor is 0.1 and the electron density 30 cm^{-3} . The observed values are represented by an horizontal line.

4.1 Description of the models

González Delgado et al. (1999b) have computed evolutionary stellar population synthesis models that predict the photospheric absorption H Balmer and HeI lines, between 3700 and 5000 Å, for a single-metallicity stellar population. The models, which are optimized for galaxies with active star formation, synthesize the profiles of the H Balmer series (H β , H γ , H δ , H8, H9, H10, H11, H12 and H13) and the

HeI absorption lines (HeI λ 4922, HeI λ 4471, HeI λ 4388, HeI λ 4144, HeI λ 4121, HeI λ 4026, HeI λ 4009 and HeI λ 3819), with a spectral sampling of 0.3 Å pixel^{-1} , for a burst and for continuous star formation at a constant rate. They use a stellar library that includes NLTE absorption profiles for stars hotter than 25000 K, and LTE profiles for lower temperatures. The temperature and gravity coverage is $4000 \leq T_{\text{eff}} \leq 50000 \text{ K}$, and $0.0 \leq \log g \leq 5.0$, respectively (González Delgado & Leitherer 1999). The models assume




Figure 8. As Figure 7, but for an instantaneous burst with a Salpeter IMF upper mass limit cut-off $120 M_{\odot}$ (filled circles), $80 M_{\odot}$ (open circles), $60 M_{\odot}$ (squares), and $30 M_{\odot}$ (open triangles).

that stars evolve from the main sequence following the evolutionary tracks of the Geneva group (Schaller et al. 1992; Schaerer et al. 1993a,b; Charbonnel et al. 1993; Meynet et al. 1994). The strength of the Balmer and HeI lines is sensitive to the age after the first 3 Myr of evolution, and sensitive to the IMF if the age is younger than 3–4 Myr. Models assume that stars have a metallicity $Z = 0.008$, and make different assumptions about the IMF slope ($\alpha = 1.5, 2.35$ and 3.0), and upper mass limit cut-off ($M_{up} = 80 M_{\odot}$ and $30 M_{\odot}$). For the purpose of this paper, the range of ages computed spans from 0 to 10 Myr.

4.2 Model Results

We use the equivalent width of HeI $\lambda 4388$, $\lambda 4026$, and $\lambda 3819$, and the wing absorption profiles of H β , H γ , H δ , H8, H9 and H10 to constrain the age, IMF and star formation law. HeI $\lambda 4388$, $\lambda 4026$, and $\lambda 3819$ are detected in absorption, because the corresponding nebular emission lines have equivalent widths that are at least a factor 3 weaker than the nebular emission in HeI $\lambda 4471$ (González Delgado et al. 1999b). Note that HeI $\lambda 4471$ is only partially filled with the nebular emission; therefore, the nebular emission of the




Figure 9. As Figure 7, but for an instantaneous burst with an IMF slope $\alpha=2.35$ (Salpeter, filled circles), $\alpha=1.5$ (open circles) and $\alpha=3.0$ (squares) and upper mass limit cut-off $80 M_{\odot}$.

lines He I $\lambda 4388$, He I $\lambda 4026$ and He I $\lambda 3819$ is $\simeq 0$, and the equivalent widths of these absorption features represent well the strength of the stellar continuum radiation of the cluster. The equivalent widths of these lines, He I $\lambda 4388$, He I $\lambda 4026$ and He I $\lambda 3819$, measured in the spectrum of Fig. 5 are 0.32, 0.55 and 0.26 Å, respectively. Their strength is compatible with continuous star formation and with burst models 3 Myr old (Fig. 10). However, He I lines can constrain the IMF. Models with an IMF steeper than Salpeter or with $M_{\text{up}} = 30 M_{\odot}$ predict He I lines stronger than ob-

served. The strengths of the lines are compatible with a 3 to 4 Myr old burst formed following a Salpeter or slightly flatter IMF ($\alpha = 1.5$, Fig. 10). The absorption wings of the Balmer lines are also compatible with burst models 3 Myr old, with Salpeter or slightly flatter IMF ($\alpha = 1.5$, Fig. 11).

burst galaxies^{||}. Most of these lines are formed in the stellar winds of the massive stars that belong to the starburst. The profile of these lines reflects the stellar massive content in the starburst; therefore, they depend on the IMF and star formation law (Leitherer, Robert & Heckman 1995). In this section, we constrain the evolutionary state and the IMF of NGC 604 by means of fitting the profiles of the ultraviolet stellar lines Si IV and C IV.

5.1 Description of the models

Evolutionary stellar population models have been computed with the code Starburst 99 (Leitherer et al. 1999). The code uses a stellar library of *IUE* ultraviolet spectra of O, B and Wolf-Rayet stars (Robert, Leitherer & Heckman 1993). The spectral resolution of the O and Wolf-Rayet stars observed is 0.75 Å. These stars are located in the solar neighborhood. However, the evolutionary models computed here assume that the stars evolve from the main sequence following the evolutionary track at $Z = 0.008$. Models are computed for instantaneous bursts between 0 and 10 Myr, and for continuous star formation lasting 10 Myr, and different assumptions about the slope ($\alpha=2.35, 3.0$ and 1.5) and upper mass limit cut-off ($M_{\text{up}} = 120, 80, 60$ and $40 M_{\odot}$) of the IMF. The model spectra are smoothed to the *IUE* spectral resolution of the observations of NGC 604, which is 6 Å.

5.2 Model results

The strongest wind stellar features in the spectra of NGC 604 are N V $\lambda 1240$, Si IV $\lambda 1400$, C IV $\lambda 1550$, He II $\lambda 1640$, and N IV $\lambda 1720$. These features are present in all the spectra of NGC 604; however, for this analysis we use the average spectrum of those at PA=149°, 136° and 159°, and the spectrum at PA=32°, because they have the best signal to noise ratio. The evolutionary state, the star formation law and the IMF in NGC 604 is constrained with the profiles of these lines.

He II shows a broad emission profile in Wolf-Rayet and O3–O5 supergiant stars; N IV shows a PCygni profile in Wolf-Rayet stars; N V and C IV show strong PCygni profiles in all O stars, and Si IV only in O supergiants. Thus, the profile of these lines in the integrated spectrum of a starburst depends strongly on the stellar content and age of the stellar cluster. In particular, Si IV shows a strong PCygni profile if the cluster formed in an instantaneous burst and its age is between 3 and 5 Myr, because within this age interval the ultraviolet light is dominated by O blue supergiants. On the other hand, the profile of C IV depends strongly on the IMF. It shows a strong PCygni profile if stars more massive than $60 M_{\odot}$ are formed in the cluster, and if the slope of the IMF is flatter than $\alpha=3.0$.

The spectrum of NGC 604 indicates that its stellar population must be dominated by massive, young O stars; thus, the stellar cluster must be young and of short duration. Continuous star formation models can be ruled out because they show Si IV weaker than observed (Fig. 12). In these models the line is diluted because the fraction of O supergiants with

respect to the total number of O stars is lower in the continuous star formation than in the instantaneous burst models. The strength of Si IV in NGC 604 indicates that it is an instantaneous burst. The age of the burst has to be between 3 to 5 Myr, because bursts younger than 3 Myr or older than 5 Myr have very few O supergiants, and thus very weak Si IV (Fig. 13). On the other hand, from the strength of C IV we can exclude instantaneous bursts with $M_{\text{up}} \leq 60 M_{\odot}$ (Fig. 14), and IMF steeper than $\alpha=3.0$ (Fig. 15). Therefore, we conclude that continuous star formation and instantaneous burst models with few very massive stars can be ruled out. The wind lines are compatible with an instantaneous burst, formed following a Salpeter or slightly flatter IMF, with upper mass limit cut-off higher than $60 M_{\odot}$. A 3 Myr instantaneous burst with Salpeter and $M_{\text{up}} \geq 80 M_{\odot}$ fits well the profile of the wind stellar lines.

6 THE MASSIVE STELLAR CONTENT OF NGC 604

We derive the massive stellar population in NGC 604 from the ultraviolet continuum luminosity, comparing the observations with the predictions of synthetic models. The number of Wolf-Rayet stars are derived from the luminosity of the He II $\lambda 1640$ line. The stellar content derived in this way is compared with that derived from *HST* photometry of stars in NGC 604. However, we need first to estimate the extinction.

6.1 Extinction estimates

Leitherer & Heckman (1995) have shown that the ultraviolet continuum arising from a young starburst (age ≤ 10 Myr) has a spectral index, β ($F_{\lambda} \propto \lambda^{\beta}$), which is independent of the IMF, star formation law and metallicity. Thus, any deviation from the predicted value, $\simeq 2.5$ for a 3 Myr instantaneous burst, could be attributed to reddening.

The ultraviolet continuum flux distribution of the *IUE* spectra of NGC 604 shows evidence of reddening, because after correcting by Galactic extinction, $E(B - V) = 0.03$ (McClure & Racine 1969), the spectra are flatter than the spectral energy distribution predicted by the evolutionary synthesis models. Massey & Hutchings (1983), analysing *IUE* spectra of H II regions in M33, conclude that the M33 extinction curve is significantly different from that of the Galaxy, because the spectra of the H II regions show very weak 2200 Å interstellar absorption dip, resembling those of the LMC and SMC. Therefore, we use the LMC curve to match the observed spectra with the spectral energy distribution of a 3 Myr old instantaneous burst. The $E(B - V)$ derived is 0.1 for the spectrum at PA=32°, and 0.12 for the average spectrum of those at PA=136°, 159° and 149°. These values are in agreement with those derived by Massey & Hutchings (1983). However, the extinction can be 0.03 higher than the values above if it is derived comparing the observed spectra with the spectral energy distribution predicted including only the stellar contribution. This is due to the nebular contribution being dominated by the two-photon continuum emission that peaks at ~ 1500 Å. If the nebular continuum is not included, the spectral energy distribution is steeper and a higher extinction is required to match the observed

^{||} One important exception is knot A in the RH II NGC 2363 (Drissen et al. 2000), so young that the effect of winds is not yet reflected in its uv spectrum.

with that derived from photometric studies based on the detection of individual stars.

ACKNOWLEDGMENTS

It is a pleasure to thank Gary Ferland and Claus Leitherer for kindly making their codes available, and Jaime Perea for the use of SIPL. Claus suggested that something should be done with those *IUE* spectra. We have benefited from stimulating and helpful discussions with members of the GEFÉ collaboration, in particular with Angeles Díaz, Jesús Maíz-Apellániz, Divakara Mayya, Guillermo Tenorio-Tagle, Elena Terlevich, Roberto Terlevich and José M. Vílchez. We are also indebted to Grazyna Stasińska for his detailed comments from a thorough reading of the paper. The *IUE* helpdesk have always been solicitous with our queries, in particular we acknowledge Pedro Rodríguez Pascual and Rosario González Riestra for support with the *IUE* spectra. *HST* images were retrieved from the ST-ECF *HST* archive.

REFERENCES

- Calzetti D., Kinney A.L., Storchi-Bergmann T., 1994, *ApJ*, 429, 582
- Cassatella A., Altamore A., González-Riestra R., Ponz J.D., Barbero J., Talavera A., Wamsteker W., 2000, *A&AS*, 141, 331
- Charbonnel C., Meynet G., Maeder A., Schaller G., Schaerer D., 1993, *A&AS*, 101, 415.
- Churchwell E., Goss W.M., 1999, *ApJ*, 514, 188
- Cid Fernandes R., Dottori H.A., Gruenwald R.B., Viegas S.M., 1992, *MNRAS*, 255, 165
- Conti P.S., Massey P., 1981, *ApJ*, 249, 471
- Clayton C.A., 1988, *MNRAS*, 231, 191
- D’Odorico S., Dopita M.A., Benvenuti P., 1980, *A&AS*, 40, 67
- D’Odorico S., Rosa M., 1981, *ApJ*, 248, 1015
- Díaz A.I., Terlevich E., Pagel B.E.J., Vílchez J.M., Edmunds M.G., 1987, *MNRAS*, 226, 19
- Drissen L., Moffat A.F.J., Shara M.M., 1990, *ApJ*, 364, 496
- , 1993, *AJ*, 105, 1400
- , Roy J.-R., Robert C., Devost D., Doyon R. 2000, *AJ*, 119, 688
- Ferland G.J., 1997, *Hazy*, a Brief Introduction to CLOUDY, University of Kentucky, Department of Physics and Astronomy Internal Report
- Freedman W.L., Wilson C.D., Madore B.F., 1991, *ApJ*, 372, 455
- García-Vargas M.L., Díaz A.I., 1994, *ApJS*, 91, 553
- , Bressan A., Díaz A.I., 1995a, *A&AS*, 112, 13
- , Bressan A., Díaz A.I., 1995b, *A&AS*, 112, 35
- , González-Delgado R.M., Pérez E., Alloin D., Díaz A.I., Terlevich E., 1997, *ApJ*, 478, 112
- González Delgado R.M., Heckman T., Leitherer C., Meurer G., Krolik J., Wilson A.S., Kinney A.L., Koratkar A., 1998, *ApJ*, 505, 174
- , García-Vargas M.L., Goldader J., Leitherer C., Pasquali A., 1999a, *ApJ*, 513, 707
- , Leitherer C., 1999, *ApJS*, 125, 479
- , Leitherer C., Heckman T., 1999b, *ApJS*, 125, 489
- González-Riestra R., Cassatella A., Solano E., Altamore A., Wamsteker W. 2000, *A&AS*, 141, 343
- Hawley S.A., Grandi S.A., 1977, *ApJ*, 217, 420
- Hippenlein H., Fried J.W., 1984, *A&A*, 141, 49
- Hunter D.A., Baum W.A., O’Neil E.J., Lynds R., 1996, *ApJ*, 456, 174
- Israel F.P., van der Kruit P.C., 1974, *A&A*, 32, 363
- Kennicutt R.C., 1984, *ApJ*, 287, 116
- Kobulnicky H.A., Zaritsky D., 1999, *ApJ*, 511, 120
- Kwitter K.B., Aller L.H., 1981, *MNRAS*, 195, 939
- Leitherer C., 1997, in Waller W.H., Fanelli M.N., Hollis J.E., Danks A.C., eds., *AIP Conf. Procs.* 408, American Institute of Physics, New York, p. 119
- Leitherer C., Heckman T.M., 1995, *ApJS*, 96, 9
- , Robert C., Drissen L. 1992, *ApJ*, 401, 596
- , Robert C., Heckman T.M., 1995, *ApJS*, 99, 173
- , Schaerer D., Goldader J.D., González Delgado R.M., Robert C., Foo Kune D., de Mello D.F., Devost D., Heckman T.M., 1999, *ApJS*, 123, 3
- , Vacca W.D., Conti P.S., Filippenko A.V., Robert C., Sargent W.L.W., 1996, *ApJ*, 465, 717
- Lejeune T., Buser R., Cuisinier F., 1997, *A&AS*, 125, 229
- Luridiana V., Peimbert M., Leitherer C., 1999, *ApJ*, 527, 110
- Madau P., Ferguson H.C., Dickinson M.E., Giavalisco M., Steidel C.C., Fruchter A., 1996, *MNRAS*, 283, 1388
- Maíz-Apellániz, J., 1999, PhD Thesis, Univ. Complutense de Madrid
- Mas-Hesse J.M., Kunth D., 1999, *A&A*, 349, 765
- Massey P., Hutchings J.B., 1983, 275, 578
- McClure R.D., Racine R., 1969, *AJ*, 74, 1000
- Medina Tanco G.A., Sabalisk N., Jatenco-Pereira V., Opher R., 1997, *ApJ*, 487, 163
- Meynet G., Maeder A., Schaller G., Schaerer D., Charbonnel C., 1994, *A&AS*, 103, 97
- Morton D.C., 1967, *ApJ*, 147, 1017
- Muñoz-Tuñón C., Tenorio Tagle G., Castañeda H.O., Terlevich R., 1996, *AJ*, 112, 1636
- Peimbert M., 1970, *PASP*, 82, 636
- Robert C., Leitherer C., Heckman T.M., 1993, *ApJ*, 418, 749
- Rodríguez-Pascual P.M., González-Riestra R., Schartel N., Wamsteker W. 1999, *A&AS*, 139, 183
- Rosa M., D’Odorico S., 1982, *A&A*, 108, 339
- Rosa M., Joubert M., Benvenuti P., 1984, *A&AS*, 57, 361
- Rosa M., Solf J., 1984, *A&A*, 130, 29
- Sabalisk N.S.P., Tenorio-Tagle G., Castañeda H.O., Muñoz-Tuñón C., 1995, *ApJ*, 444, 200
- Schaerer D., Charbonnel C., Meynet G., Maeder A., Schaller G., 1993a, *A&AS*, 102, 339
- Schaerer D., Meynet G., Maeder A., Schaller G., 1993b, *A&AS*, 98, 523
- Schaller G., Schaerer D., Meynet G., Maeder A., 1992, *A&AS*, 96, 269
- Schmutz W., Leitherer C., Gruenwald R.B., 1992, *PASP*, 104, 1164
- Smith H.E., 1975, *ApJ*, 199, 591
- Stasińska G., Leitherer C., 1996, *ApJS*, 107, 661
- Stasińska G., Schaerer D., 1999, *A&A*, 351, 72
- Terlevich E., Díaz A.I., Terlevich R., González Delgado R.M., Pérez E., García-Vargas M.L., 1996, *MNRAS*, 279, 1219
- Vacca W.D., Robert C., Leitherer C., Conti P.S., 1995, *ApJ*, 444, 647
- Viallefond F., Goss W.M., 1986, *A&A*, 154, 357
- Vílchez J.M., Pagel B.E.J., 1988, *MNRAS*, 231, 257
- Walborn N.R., 1991, in Leitherer C., Walborn N., Heckman T., Norman C., eds, *Massive Stars in Starbursts*. Cambridge Univ. Press, Cambridge, p. 145
- Walborn N.R., Fitzpatrick E. L., 1990, *PASP*, 102, 379
- Yang H., Chu Y.-H., Skillman E.D., Terlevich R., 1996, *AJ*, 112, 146

This figure "fig1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0003067v1>

This figure "fig2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0003067v1>